

Unconventional temperature-enhanced magnetism in $\text{Fe}_{1.1}\text{Te}$

I. A. Zaliznyak^a, Z. J. Xu^a, J. M. Tranquada^a, G. D. Gu^a, A. M. Tsvelik^a, and M. B. Stone^b

^aCondensed Matter Physics & Materials Science Dept., Brookhaven National Lab, Upton, NY, USA

^bOak Ridge National Lab, Oak Ridge, TN, USA

There are two common scenarios used to describe the magnetism in the families of Fe-based superconductors. In one, the magnetism originates from local atomic spins, while in the other it corresponds to a cooperative spin-density-wave (SDW) behavior of conduction electrons. Both assume clear partition into localized electrons, giving rise to local spins, and itinerant ones, occupying well-defined, rigid conduction bands. We have used inelastic neutron scattering to characterize both the static and dynamic magnetism in a crystal of $\text{Fe}_{1.1}\text{Te}$, parent to the $\text{Fe}_{1+y}\text{Te}_{1-x}\text{Se}_x$ family of superconductors.¹ In contrast to the simple pictures, we find that localized spins and itinerant electrons are coupled together. In particular, we have evaluated the effective magnetic moment by integrating both the elastic and inelastic magnetic scattering. The effective spin per Fe at $T \approx 10$ K, in the antiferromagnetic phase, corresponds to $S \approx 1$, consistent with the recent analyses that emphasize importance of Hund's intra-atomic exchange. However, it grows to $S \approx 3/2$ in the disordered phase, a result that presents a challenge to current theoretical models.

Work at Brookhaven is supported by the Office of Basic Energy Sciences, Division of Materials Science and Engineering, U.S. Department of Energy, under Contract No. DE-AC02-98CH10886.

1. I. A. Zaliznyak *et al.*, arXiv:1103.5073.