

Angewandte International Edition www.angewandte.org

High-Pressure Chemistry

How to cite: Angew. Chem. Int. Ed. 2025, e202422710 doi.org/10.1002/anie.202422710

Pressure-Driven Reactivity in Dense Methane-Nitrogen Mixtures

Hannah A. Shuttleworth, Mikhail A. Kuzovnikov, Lewis J. Conway, Huixin Hu, Jinwei Yan, Samuel Gallego-Parra, Israel Osmond, Tomas Marqueño, Michael Hanfland, Dominique Laniel, Eugene Gregoryanz, Andreas Hermann,* Miriam Peña-Alvarez,* and Ross T. Howie*

Abstract: Carbon, nitrogen, and hydrogen are among the most abundant elements in the solar system, and our understanding of their interactions is fundamental to prebiotic chemistry. CH_4 and N_2 are the simplest archetypical molecules formed by these elements and are both markedly stable under extremes of pressure. Through a series of diamond anvil cell experiments supported by density functional theory calculations, we observe diverse compound formation and reactivity in the CH_4 - N_2 binary system at high pressure. Above 7 GPa two concentration-dependent molecular compounds emerge, $(CH_4)_5N_2$ and $(CH_4)_7(N_2)_8$, held together by weak van der Waals interactions. Strikingly, further compression at room temperature irreversibly breaks the N_2 triple bond, inducing the dissociation of CH_4 above 140 GPa, with the near-quenched samples revealing distinct spectroscopic signatures of strong covalently bonded C-N-H networks. High temperatures vastly reduce the required pressure to promote the reactivity between CH_4 and N_2 , with NH_3 forming together with longer-chain hydrocarbons at 14 GPa and 670 K, further decomposing into powdered diamond when temperatures exceed 1200 K. These results exemplify how pressure-driven chemistry can cause unexpected complexity in the most simple molecular precursors.

Introduction

Carbon, nitrogen, and hydrogen constitute some of the most prevalent elements in the solar system, and exploring their

[*] H. A. Shuttleworth, M. A. Kuzovnikov, L. J. Conway, J. Yan,
I. Osmond, T. Marqueño, D. Laniel, E. Gregoryanz, A. Hermann,
M. Peña-Alvarez, R. T. Howie
Centre for Science at Extreme Conditions, University of Edinburgh,
Edinburgh, EH9 3FD, United Kingdom
E-mail: ross.howie@ed.ac.uk
Miriam.Pena.Alvarez@ed.ac.uk
a.hermann@ed.ac.uk
H. Hu, R. T. Howie
Center for High Pressure Science and Technology Advanced
Research, 1690 Cailun Road, Shanghai, 201203, China
L. I. Conway

Department of Materials Science & Metallurgy, University of Cambridge, 27 Charles Babbage Road, Cambridge CB3 0FS, United Kingdom

S. Gallego-Parra, M. Hanfland

European Synchrotron Radiation Facility, 71 Avenue des Martyrs, 38000 Grenoble, France

E. Gregoryanz

Key Laboratory of Materials Physics, Institute of Solid State Physics, CAS, Hefei, 230031, China

E. Gregoryanz

SHARPS (Shanghai Advanced Research in Physical Sciences), 68 Huatuo Road, Shanghai, 201203, China

© 2025 The Author(s). Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited. chemical interactions is crucial to understanding the origin of life.^[1] To explore the chemical interplay between these elements at planetary relevant conditions, the effect of pressure must be considered in addition to temperatureinduced chemistry. By altering molecular orbitals and interactions, high pressure serves as a clean chemical method for the synthesis of novel materials that would otherwise be unstable at ambient pressure.^[2-12] A key example of this is how the stability and reactivity of molecular materials evolve upon compression. Due to the extremely strong N \equiv N triple bond, molecular N₂ is one of the least reactive molecules under ambient conditions. However, high pressure has been shown to induce the formation of numerous non-molecular phases of nitrogen, requiring the breaking of the triple bond.^[13-22] The potential chemistry of N₂ at high pressure is further enhanced with temperature, resulting in the synthesis of novel polynitrides, which are of key interest due to their potential as high energy density compounds.^[4,10,23–28]

Like nitrogen, methane is another example of a simple molecular system that undergoes a series of physical and structural changes under pressure, with crucial implications in understanding physical processes within planetary interiors.^[29–31] Both theoretical and experimental studies have demonstrated that the decomposition of methane into longer hydrocarbons and diamond is favored when compressed and heated.^[29,32–38] Additionally, it is theorized that at room temperature and pressures approaching 300 GPa, methane will dissociate into diamond and H₂, which is of great interest regarding the mantle region of planets such as Neptune and Uranus, where methane is abundant.^[33,39–42]

Angew. Chem. Int. Ed. 2025, e202422710 (1 of 7)

There is observational evidence that reactions occur between CH₄ and N₂ in the atmosphere of the satellite Titan, whereby hydrocarbon-nitrile aerosol compounds, named "tholins", have formed from CH₄-N₂ mixtures at low pressures and temperatures in the presence of solar ultraviolet radiation.^[43-47] Recent theoretical works also predict that under dense planetary conditions, exotic organic chemistry occurs within the C-N-H system.^[48,49] A variety of stable C-N-H compounds are predicted to emerge at pressures up to 50 GPa.^[48] Most notably, two high energy density compounds were predicted: CN₂H₄ and CH₄N₄, stable at 11 GPa and 41 GPa respectively, the latter of which is predicted to be stable to ambient conditions upon quenching. With a calculated energy release of 6.43 KJg⁻¹ upon decomposition back into molecular N2 and CH4, this is a promising high energy density compound.^[48]

At ambient pressure, experiments have shown that CH_4 and N_2 can react, though require combinations of hightemperature, electric discharge, microwave radiation, and catalysts.^[50-53] Most experimental approaches investigating the C–N–H system at high pressure have used complex precursors rather than the simplest archetypical molecules originating from the ternary system: CH_4 and N_2 .^[54-57] There has been only one experimental study on CH_4 - N_2 mixtures up to 16 GPa, presenting evidence of compound formation retaining N_2 and CH_4 molecular units; however, the structural characterization was unsuccessful.^[58] Surprisingly, whether the combination of high pressure and temperature can facilitate a reaction between CH_4 and N_2 has yet to be explored experimentally.

In this work, we investigate pressure and temperatureinduced chemistry in the dense CH_4 - N_2 system. We report the formation of two CH_4 - N_2 van der Waals compounds above 7 GPa at room temperature from binary CH_4 - N_2 fluid mixtures. Through a combination of X-ray diffraction and first principle calculations, these are identified as $P4_2/mnm$ - $(CH_4)_7(N_2)_8$ and *Ibam*- $(CH_4)_5N_2$. Remarkably, upon room temperature compression of either compound, we observe the irreversible pressure-induced dissociation and the reaction of N_2 and CH_4 molecules above 140 GPa. Decompression of the reaction products to near-ambient conditions reveals spectroscopic signatures of strong covalently bonded C–N–H networks. The application of high temperature can induce a reaction at substantially lower pressures, where we observe the formation of NH₃, together with longer-chain hydrocarbons at 670 K (at 14 GPa), the latter of which decomposes into powdered diamond when temperatures exceed 1200 K at 25 GPa.

Results and Discussion

Synthesis, Characterization and Stability of $\mathsf{CH}_4\text{-}\mathsf{N}_2$ Molecular Compounds

Two representative CH_4 - N_2 gas mixtures (50:50 CH_4 : N_2 and 67:33 CH_4 : N_2) were loaded into diamond anvil cells at 0.2 GPa (see Supporting Information for a complete description of the experimental procedure). Upon compression above 2 GPa, the fluid mixtures solidified into A- CH_4 + δ - N_2 .^[59-63] Compression of the 50:50 mixture above 7 GPa resulted in the formation of a new compound. Single crystal X-ray diffraction (SCXRD) data were obtained. These datasets were successfully refined to a crystal structure isomorphic to σ -CrFe (space group $P4_2/mnm$, Figure 1a), with lattice parameters a=11.904(3) Å and c=6.2072(13) Å,



Figure 1. (a) Structural models of the unit cell of $P4_2/mnm$ -(CH₄)₇(N₂)₈ (top) and a projection of the $P4_2/mnm$ -(CH₄)₇(N₂)₈ structure along the *c*-axis (bottom). Different colors represent different molecular sites rather than different molecular types. The shape of the ellipsoids represents the preferred orientation of the molecules. (b) Slice of the (0kl) reciprocal space of an $P4_2/mnm$ -(CH₄)₇(N₂)₈ single crystal at 7 GPa. The blue circles indicate visible reflections. The systematic absences are consistent with those of space group $P4_2/mnm$. (c) Reconstruction of the (hk2) reciprocal space slice built using single-crystal data collected on *Ibam*-(CH₄)₅N₂ single crystal at 13 GPa. The blue circles highlight visible reflections. The systematic absences are consistent with cell of *Ibam*-(CH₄)₅N₂ (top) and a projection of the *Ibam*-(CH₄)₅N₂ unit cell along the *c*-axis (bottom), nitrogen atoms are represented as blue spheres and CH₄ molecules are represented by brown spheres.

Angew. Chem. Int. Ed. 2025, e202422710 (2 of 7)

at 7 GPa (see Figure 1b and Supporting Table S1 for crystallographic data).^[64] The unit cell of this compound is shown in Figure 1a. Due to strong molecular rotational disorder providing limited reflections at low *d*-spacings, along with a comparable scattering factor for N_2 and CH₄, we were unable to estimate site occupancies. We observed no further structural changes in our powder XRD measurements (Supporting Figure S1a) up to at least 47 GPa (Figure 2b).

Upon compression of the 67:33 CH₄:N₂ mixture to 9 GPa, we observe the presence of another uncharacterized compound. SCXRD data obtained at 13 GPa were refined to an *Ibam* unit cell (Figure 1d) with lattice parameters a =11.854(12) Å, b=18.459(5) Å and c=5.6278(12) Å at 13 GPa (see Figure 1c and Supporting Table S2 for crystallographic data).^[68] To the best of our knowledge, this is the first observation of this structural type. Out of the six distinct crystallographic positions, nitrogen could unambiguously be assigned to the 16k site (see Supporting Information for further details). With the five remaining sites assigned as CH₄ molecules, the experimental data suggests a (CH₄)₅N₂ composition. Extracted volumetric data from powder XRD measurements (Supporting Figure S1b) upon compression demonstrates that this compound is stable to at least 23 GPa (Figure 2b).

To ascertain the stability of the molecular compounds and estimate their compositions, we performed density functional theory (DFT) calculations (see Supporting Information for computational details). The resulting lowest formation enthalpies form part of Figure 2a, where they are shown relative to pure ε -N₂ and pure CH₄ (in the HP phase, *R*3-CH₄).^[66] The convex hull also includes additional data points based on the δ -N₂ structure type, which was proposed in earlier work^[58] to allow for some uptake of CH₄. The only compositions that emerge as stable phases on the convex hull are $P4_2/mnm$ -(CH₄)₇(N₂)₈ and *Ibam*-(CH₄)₅N₂, the latter agreeing with experimental results. The Supporting Information contains molecular dynamics (MD) simulation results (which confirm the compounds' kinetic stability and molecular rotational/librational character), composition-volume data, and results from alternative exchange-correlation functionals.

We have investigated both the $P4_2/mnm$ -(CH₄)₇(N₂)₈ and Ibam-(CH₄)₅N₂ phases using Raman spectroscopy, which allows us to directly explore any changes in the inter/intramolecular environments. At 8.5 GPa, the Raman spectrum of $P4_2/mnm$ -(CH₄)₇(N₂)₈ exhibits the characteristic Raman modes of molecular CH₄ and N₂: the C–H bending mode ν_2 at 1536 cm⁻¹; C–H symmetric (ν_1) and antisymmetric (ν_3) modes at 2996 cm⁻¹ and 3115 cm⁻¹, respectively; and the N₂ intramolecular vibrational modes, ν_2 at 2341 cm⁻¹ and ν_1 at 2346 cm⁻¹. At lower pressures, the Raman spectra of Ibam-(CH₄)₅N₂ closely resembles that of P4₂/mnm-(CH₄)₇- $(N_2)_8$. Upon compression, it becomes evident that the Raman frequency of the N₂ vibrons and the CH₄ stretching modes of each compound have a different pressure dependency, indicative that the molecular environments differ. For example, at 28 GPa, the Raman frequency of the N₂- ν_2 vibron is 2372 cm⁻¹ for $P4_2/mnm$ -(CH₄)₇(N₂)₈, and 2367 cm⁻¹ for *Ibam*-(CH_4)₅N₂.

Additionally, both compounds exhibit a distinct deviation in the frequencies of the Raman active modes compared with the pure species (Figure 3a).^[31,67] For example, in pure nitrogen, there is complex splitting of the N₂- ν_2 vibron at 24 GPa^[67] (Supporting Figure S2), whilst P4₂/mnm-(CH₄)₇-(N₂)₈ and *Ibam*-(CH₄)₅N₂ show two distinct modes. Splitting of the CH₄- ν_1 into CH₄- $\nu_1(1)$ and CH₄- $\nu_1(2)$ is observed during compression of *Ibam*-(CH₄)₅N₂ beyond 67 GPa^[31]



Figure 2. (a) Ground state relative enthalpies of CH_4 - N_2 mixtures from DFT calculations at 15 GPa, based on the σ -CrFe, *Ibam*, and δ - N_2 structure types (see text). The black line indicates the convex hull of stable phases. (b) Volume per molecule of the two van der Waals compounds as a function of pressure, obtained from powder XRD data. Symbols represent experimental data for $P4_2/mnm$ -(CH_4)₇(N_2)₈ (blue) and *Ibam*-(CH_4)₅ N_2 (orange), with stars representing SCXRD data and the remaining symbols representing powder XRD data. The solid lines represent their best Birch-Murnaghan fit. The dashed line represents the equation of state of pure N_2 ⁽⁶⁵⁾ and the dotted line represents that of pure CH_4 .

Angew. Chem. Int. Ed. 2025, e202422710 (3 of 7)



Figure 3. (a) Raman spectra upon compression of $P4_2/mnm$ -(CH₄)₇(N₂)₈, synthesized from a 50% N₂ mixture, from 8.5 to 163 GPa (blue), and the Raman spectrum of pure CH₄ at 163 GPa (lilac). The asterisk denotes a peak which is an artifact of the diamond anvils. (b) Raman shift of the N₂ vibrational modes and C–H stretching modes of $P4_2/mnm$ -(CH₄)₇(N₂)₈ (blue) and *Ibam*-(CH₄)₅N₂ (orange) as a function of pressure, synthesized from a 50% and 67% CH₄ gas mixture, respectively. Different symbols represent different experimental runs, and the empty symbols denote data collected upon decompression. Dotted lines represent the N₂ vibrational modes of pure CH₄.^[31] Inset: Profile of the N₂ vibrational modes of $P4_2/mnm$ -(CH₄)₇(N₂)₈ (blue) and *Ibam*-(CH₄)₅N₂ (orange) at 30 GPa.

(Figure 3b). The splitting is not resolvable in $P4_2/mnm$ -(CH₄)₇(N₂)₈ due to the low intensity of the C–H stretching modes. Furthermore, inspection of the N₂ vibrational modes reveals that the relative intensity of the N₂- ν_1 and N₂- ν_2 modes also depends on the compound formed, as seen in Figure 3b. In both compounds, we also see an intense broad band at frequencies close to the Rayleigh line (Figure 3a). This could be attributed to rotational disordered molecules, which is in agreement with our X-ray diffraction measurements and MD simulations demonstrating freely rotating CH₄ units (Supporting Figure S6).

Pressure Induced Reactivity in the CH₄-N₂ System

While CH₄ is stable to at least 200 GPa,^[31,69] the triple bond of molecular N₂ undergoes progressive weakening to the extent that polymeric allotropes form above 110 GPa.^[13-22] We investigated the reactivity of $P4_2/mnm$ -(CH₄)₇(N₂)₈ (Figure 3a) and *Ibam*-(CH₄)₅N₂ (Supporting Figure S3) phases using Raman spectroscopy. Compression of $P4_2/mnm$ -(CH₄)₇(N₂)₈ led to a turnover in the frequencies of the N₂ vibrons above 100 GPa and subsequent softening of the modes (Figure 3b). At 128 GPa, the higher frequency ν_1 vibron becomes unidentifiable, while the N₂- ν_2 mode is unresolvable above 163 GPa in $P4_2/mnm$ -(CH₄)₇(N₂)₈ and 136 GPa in *Ibam*-(CH₄)₅N₂. Before disappearing, the frequency of the N₂ vibrational mode in $P4_2/mnm$ -(CH₄)₇(N₂)₈ at 163 GPa is comparative to the frequency observed in pure N₂ at the onset of polymerization (120 GPa).^[16,19] This is indicative of pressure-induced breaking of the N₂ triple bond to form polymeric nitrogen.^[13,14,16,67,70-73] Concurrently, a new mode at around 1200 cm⁻¹ appears, corresponding to vibrational modes of single bonded C–N and C–C.^[74-77] Additionally, the CH₄- ν_2 bending mode, along with the symmetric (ν_1) and anti-symmetric (ν_3) stretching modes, decreases in intensity and broadens (Figure 3a). A direct comparison to the Raman spectrum of CH₄,^[31] which we measured at 163 GPa to provide a reference (Figure 3a), indicates dissociation and formation of a reaction product. No detectable X-ray diffraction patterns were obtained at this pressure, likely due to weak scattering or sample amorphization, similar to η -nitrogen.^[14]

Strikingly, decompression of the reaction products to 5(4) GPa, does not result in the transformation back into either molecular compound nor do the Raman signatures of molecular N₂ reappear (Figure 4a and Supporting Figure S3), unlike pure N₂ whereby the vibrons re-emerge at 40 GPa.^[67] Similarly, the C–H stretching modes remain broad compared to either molecular compound at comparative pressures, while the band between 1100–1200 cm⁻¹ remains active but as a broad series of peaks, which we attribute to the formation of oligomers.^[74] These Raman features are observed regardless of whether the reaction products of $P4_2/mnm$ -(CH₄)₇(N₂)₈ or *Ibam*-(CH₄)₅N₂ are decompressed.

By comparing the experimental data with the known Raman spectra of the simplest covalent C–H–N compounds, amines, we tentatively assign it to propylamine (CH₃ (CH₂)₂NH₂). As such, we measured the Raman spectra of

Angew. Chem. Int. Ed. 2025, e202422710 (4 of 7)



Figure 4. (a) Raman spectrum of NH₃, powdered diamond and hydrocarbons at 25 GPa, formed after laser heating $P4_2/mnm$ -(CH₄)₇(N₂)₈ (red), the Raman spectrum after compression of a 50% CH₄ mixture to 163 GPa then decompression to 5 GPa (black), and the Raman spectrum of propylamine (CH₃)₂(NH)₂) at 3 GPa (green). The diamond symbols represent the second-order diamond Raman mode. (b) Rietveld refinement of *fcc* NH₃-III and powdered diamond quenched to ambient temperature, formed after laser heating $P4_2/mnm$ -(CH₄)₇(N₂)₈ to above 1200 K at 40 GPa. The diffraction pattern of $P4_2/mnm$ -(CH₄)₇(N₂)₈ before laser heating is also shown.

propylamine up to 3 GPa to provide a direct comparison (Figure 4). Here, we see reasonable agreement between the modes around 3100-3450 cm⁻¹, attributed to N-H stretching, together with C-H stretching modes in the 2800-3000 cm⁻¹ region.^[76] Deformation modes within the 1400- 1500 cm^{-1} region and C-H bending in the $1560-1580 \text{ cm}^{-1}$ region are in good agreement with those of our products.^[76-78] We also see an overlap in the N–H stretching region, indicating the presence of additional molecular units. Unfortunately, attempts at structural characterization of the reaction products were unsuccessful, which we attribute to the small sample sizes combined with the weak X-ray scattering of the constituent elements, whilst full decompression led to the loss of the samples. In recent theoretical works exploring compound formation in the C-N-H system, a molecular compound composed of propylamine, NH₃, and CH₄ units is predicted to be the most thermodynamically stable reaction product in CH₄-N₂ mixtures below 10 GPa.^[48] As such, we propose this molecular compound, CH₃ $(CH_2)_2NH_2 + NH_3 + CH_4$, as the most promising candidate for the compression product. Interestingly, amines, including propylamine, are believed to be a constituent of Titan's "tholins".[79,80]

Given that temperature is known to destabilize both CH_4 and N_2 at high pressure, we explored the high temperature stability of the molecular compounds in an attempt to synthesize covalently bonded C–N–H compounds at lower pressures.^[48] Upon heating *Ibam*-(CH₄)₅N₂ phase to temperatures above 670 K at 14 GPa, we observe a reaction to form NH₃ as evidenced by our X-ray diffraction measurements (see Supporting Figure S4).^[81] Laser heating either *P*4₂/ *mnm*-(CH₄)₇(N₂)₈ or *Ibam*-(CH₄)₅N₂ precursors at 40 GPa to temperatures in excess of 1200 K also led to a reaction, with the temperature quenched diffraction pattern showing the formation of NH₃-III together with *fcc* diamond (*a*= 3.4778 Å) (Figure 4b and Supporting Figure S5). The presence of powdered diamond is a result of the decomposition of CH₄ into C, with H₂ reacting to form NH₃ and longer chain hydrocarbons when laser heated.^[29,33,34] Raman spectroscopy experiments of the quenched sample support these observations, with the N₂ vibrons becoming unresolvable and the NH₃ stretching modes, v_1 , $2v_4$ and v_3 , appearing at 3187 cm^{-1} , 3318 cm^{-1} and 3395 cm^{-1} at 25 GPa. The C–H stretching mode v_1 , additional modes in the C–H bending region 1460–1800 cm⁻¹, and a complex spectrum below 1330 cm^{-1} suggest the presence of a mixture of longer-chain light hydrocarbons.^[36,82] Such mixtures have been previously established as oligomers, which are undetectable in X-ray diffraction experiments.^[83]

Angewandte

Chemie

Conclusion

By compressing mixtures of the simple CH₄-N₂ binary molecular system in a series of diamond anvil cell experiments, we report the formation of two molecular van der *Ibam*-(CH_4)₅ N_2 Waals compounds, and $P4_{2}/mnm$ - $(CH_4)_7(N_2)_8$, above 7 GPa and 300 K. Remarkably, compression of either of these compounds results in the breaking of the N₂ triple bond and dissociation of CH₄, with the reaction product exhibiting spectroscopic signatures of singly-bonded C-N-H compounds upon decompression to near-ambient pressure. We identify the previously predicted molecular compound $CH_3(CH_2)_2NH_2 + NH_3 + CH_4$ as a potential reaction product. Upon heating either Ibam-(CH₄)₅N₂ or $P4_2/mnm$ -(CH₄)₇(N₂)₈ to temperatures of 670 K at 14 GPa, we observe decomposition into NH₃ and longer-chain light hydrocarbons. Above 1200 K at pressures exceeding 25 GPa, the hydrocarbons further decompose, producing powdered diamond. These combined results demonstrate the complexity of compound formation in the C-N-H ternary system under planetary-relevant conditions, even when starting from the simplest precursors. Such mixtures comprise the atmosphere of Saturn's moon Titan, and it will be of great interest to see if the reaction products we observe in experiments will be observed directly through the proposed Dragonfly mission exploring prebiotic chemistry on the satellite.

Acknowledgements

Dr. Ross T. Howie acknowledges that the project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Program (Grant Agreement 948895 "MetElOne"). Dr. Miriam Peña-Alvarez and Dr. Dominique Laniel acknowledge the support of the UK Research and Innovation (UKRI) Future Leaders Fellowship (Mrc-Mr/ T043733/1 and MR/V025724/1, respectively). The authors thank Dr. Mengan Wang for assistance in data analysis, and Ms Zena Younes for useful discussions. We acknowledge the European Synchrotron Radiation Facility (ESRF) for the provision of synchrotron radiation facilities at beamline ID15B under proposal HC-5261. We acknowledge DESY (Hamburg, Germany), a member of the Helmholtz Association HGF, for the provision of experimental facilities. Parts of this research were carried out at PETRA-III P02.2 and we would like to thank Dr. Konstantin Glazyrin for assistance. Beamtime was allocated for proposal I-20221366. Parts of the synchrotron radiation experiments were performed at BL10XU of SPring-8 with the approval of the Japan Synchrotron Radiation Research Institute (JASRI) (Proposal No. 2023B1420). Part of the study was performed at I15, Diamond Light Source (Didcot, UK) under proposal CY30698-1. We are grateful for computational support from the UK national high-performance computing service, ARCHER2, for which access was obtained via the UKCP consortium and funded by EPSRC grant ref EP/X035891/1. For open access, the author has applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising from this submission.

Conflict of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Keywords: high-pressure chemistry · raman spectroscopy · synchrotron x-ray diffraction · molecular systems · high-temperature chemistry

Angew. Chem. Int. Ed. 2025, e202422710 (6 of 7)

- [2] W. Vos, L. Finger, R. Hemley, J. Hu, H. Mao, J. Schouten, Nature 1992, 358, 46.
- [3] P. Loubeyre, M. Jean-Louis, R. LeToullec, L. Charon-Gérard, Phys. Rev. Lett. 1993, 70, 178.
- [4] E. Gregoryanz, C. Sanloup, M. Somayazulu, J. Badro, G. Fiquet, R. Hemley, Nat. Mater. 2004, 3, 294.
- [5] M. Somayazulu, P. Dera, A. Goncharov, S. Gramsch, H.-P. Liermann, Z. Liu, R. Hemley, Nat. Chem. 2010, 2, 50.
- [6] D.K. Spaulding, G. Weck, P. Loubeyre, F. Datchi, P. Dumas, M. Hanfland, Nat. Commun. 2014, 5, 5739.
- [7] A. Dewaele, N. Worth, C. J. Pickard, R. J. Needs, S. Pascarelli, O. Mathon, M. Mezouar, T. Irifune, Nat. Chem. 2016, 8, 784.
- [8] R.T. Howie, R. Turnbull, J. Binns, M. Frost, P. Dalladay-Simpson, E. Gregoryanz, Sci. Rep. 2016, 6, 34896.
- [9] A. Drozdov, P. Kong, V. Minkov, S. Besedin, M. Kuzovnikov, S. Mozaffari, L. Balicas, F. Balakirev, D. Graf, V. Prakapenka, et al., Nature 2019, 569, 528.
- [10] Y. Wang, M. Bykov, I. Chepkasov, A. Samtsevich, E. Bykova, X. Zhang, S.-q. Jiang, E. Greenberg, S. Chariton, V.B. Prakapenka, et al., Nat. Chem. 2022, 14, 794.
- [11] U. Ranieri, L. J. Conway, M.-E. Donnelly, H. Hu, M. Wang, P. Dalladay-Simpson, M. Peña-Alvarez, E. Gregoryanz, A. Hermann, R. T. Howie, Phys. Rev. Lett. 2022, 128, 215702.
- [12] J. C. Crowhurst, A. F. Goncharov, B. Sadigh, C. L. Evans, P. G. Morrall, J. L. Ferreira, A. Nelson, Science 2006, 311, 1275.
- [13] A.F. Goncharov, E. Gregoryanz, H.-k. Mao, Z. Liu, R.J. Hemley, Phys. Rev. Lett. 2000, 85, 1262.
- [14] E. Gregoryanz, A. F. Goncharov, R. J. Hemley, H.-k. Mao, Phys. Rev. B 2001, 64, 052103.
- [15] E. Gregoryanz, A. F. Goncharov, R. J. Hemley, H.-k. Mao, M. Somayazulu, G. Shen, Phys. Rev. B 2002, 66, 224108.
- [16] M. Eremets, A. Gavriliuk, I. Troyan, D. Dzivenko, R. Boehler, Nat. Mater. 2004, 3, 558.
- [17] E. Gregoryanz, A. F. Goncharov, C. Sanloup, M. Somayazulu, H.-k. Mao, R. J. Hemley, J. Chem. Phys. 2007, 126, 184505.
- C. Ji, A. A. Adeleke, L. Yang, B. Wan, H. Gou, Y. Yao, B. Li, [18] Y. Meng, J. S. Smith, V. B. Prakapenka, et al., Sci. Adv. 2020, 6. eaba9206.
- [19] D. Laniel, B. Winkler, T. Fedotenko, A. Pakhomova, S. Chariton, V. Milman, V. Prakapenka, L. Dubrovinsky, N. Dubrovinskaia, Phys. Rev. Lett. 2020, 124, 216001.
- [20] D. Laniel, G. Geneste, G. Weck, M. Mezouar, P. Loubeyre, Phys. Rev. Lett. 2019, 122, 066001.
- [21] L. Lei, Q.-Q. Tang, F. Zhang, S. Liu, B.-B. Wu, C.-Y. Zhou, Chin. Phys. Lett. 2020, 37, 068101.
- [22] A.F. Goncharov, I.G. Batyrev, E. Bykova, L. Brüning, H. Chen, M. F. Mahmood, A. Steele, N. Giordano, T. Fedotenko, M. Bykov, Phys. Rev. B 2024, 109, 064109.
- [23] D. Laniel, B. Winkler, E. Koemets, T. Fedotenko, M. Bykov, E. Bykova, L. Dubrovinsky, N. Dubrovinskaia, Nat. Commun. 2019, 10, 4515.
- [24] M. Bykov, E. Bykova, G. Aprilis, K. Glazyrin, E. Koemets, I. Chuvashova, I. Kupenko, C. McCammon, M. Mezouar, V. Prakapenka, et al., Nat. Commun. 2018, 9, 2756.
- [25] D. Laniel, G. Weck, G. Gaiffe, G. Garbarino, P. Loubeyre, J. Phys. Chem. Lett. 2018, 9, 1600.
- [26] B. A. Steele, E. Stavrou, J. C. Crowhurst, J. M. Zaug, V. B. Prakapenka, I. I. Oleynik, Chem. Mater. 2017, 29, 735.
- [27] D. Laniel, G. Weck, P. Loubeyre, Inorg. Chem. 2018, 57, 10685.
- [28] A.F. Young, C. Sanloup, E. Gregoryanz, S. Scandolo, R.J. Hemley, H.-k. Mao, Phys. Rev. Lett. 2006, 96, 155501.
- [29] L. J. Conway, A. Hermann, Geosciences 2019, 9, 227.
- [30] H. Maynard-Casely, C. Bull, M. Guthrie, I. Loa, M. McMahon, E. Gregoryanz, R. Nelmes, J. Loveday, J. Chem. Phys. 2010, 133.064504.



^[1] J. Oró, S. Kamat, Nature 1961, 190, 442.

5213773, 0 Downloaded from https://oninelibrary.wiley.com/doi/10.1002/anie.202422710 by University OF Ednburgh, Wiley Online Library on [01/05/2025]. See the Terms and Conditions (https://oninelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

- [31] J. Proctor, H. Maynard-Casely, M. Hakeem, D. Cantiah, J. Raman Spectrosc. 2017, 48, 1777.
- [32] M. Frost, R. S. McWilliams, E. Bykova, M. Bykov, R. J. Husband, L. M. Andriambariarijaona, S. Khandarkhaeva, B. Massani, K. Appel, C. Baehtz, et al., *Nature Astronomy* 2024, 8, 174–181.
- [33] L. R. Benedetti, J. H. Nguyen, W. A. Caldwell, H. Liu, M. Kruger, R. Jeanloz, *Science* 1999, 286, 100.
- [34] H. Hirai, K. Konagai, T. Kawamura, Y. Yamamoto, T. Yagi, *Phys. Earth Planet. Inter.* 2009, 174, 242.
- [35] B. Cheng, S. Hamel, M. Bethkenhagen, Nat. Commun. 2023, 14, 1104.
- [36] A. Kolesnikov, V. G. Kutcherov, A. F. Goncharov, *Nat. Geosci.* 2009, 2, 566.
- [37] L. Spanu, D. Donadio, D. Hohl, E. Schwegler, G. Galli, Proc. Natl. Acad. Sci. USA 2011, 108, 6843.
- [38] A. S. Naumova, S. V. Lepeshkin, A. R. Oganov, J. Phys. Chem. C 2019, 123, 20497.
- [39] M. Ross, Nature 1981, 292, 435.
- [40] M. A. Neumann, W. Press, C. Nöldeke, B. Asmussen, M. Prager, R. M. Ibberson, J. Chem. Phys. 2003, 119, 1586.
- [41] G. Gao, A. R. Oganov, Y. Ma, H. Wang, P. Li, Y. Li, T. Iitaka, G. Zou, J. Chem. Phys. 2010, 133, 144508.
- [42] D. Kraus, J. Vorberger, A. Pak, N. Hartley, L. Fletcher, S. Frydrych, E. Galtier, E. Gamboa, D. O. Gericke, S. Glenzer, et al., *Nature Astronomy* **2017**, *1*, 606.
- [43] J. H. Waite, D. T. Young, T. E. Cravens, A. J. Coates, F. J. Crary, B. Magee, J. Westlake, *Science* 2007, *316*, 870.
- [44] C. Sagan, B. Khare, Nature 1979, 277, 102.
- [45] J. H. Waite Jr, H. Niemann, R. V. Yelle, W. T. Kasprzak, T. E. Cravens, J. G. Luhmann, R. L. McNutt, W.-H. Ip, D. Gell, V. De La Haye, et al., *Science* 2005, *308*, 982.
- [46] C. McKay, A. Coustenis, R. Samuelson, M. Lemmon, R. Lorenz, M. Cabane, P. Rannou, P. Drossart, *Planet. Space Sci.* 2001, 49, 79.
- [47] A. Coustenis, R. K. Achterberg, B. J. Conrath, D. E. Jennings, A. Marten, D. Gautier, C. A. Nixon, F. M. Flasar, N. A. Teanby, B. Bézard, et al., *Icarus* 2007, 189, 35.
- [48] F. Peng, Y. Ma, A. Hermann, M. Miao, Phys. Rev. Mater. 2020, 4, 103610.
- [49] A. S. Naumova, S. V. Lepeshkin, P. V. Bushlanov, A. R. Oganov, J. Phys. Chem. A 2021, 125, 3936.
- [50] N. Ning, C. Qian, S. Zhou, Catalysis Science, Technology 2025, 15, 249.
- [51] X. Bai, S. Tiwari, B. Robinson, C. Killmer, L. Li, J. Hu, *Catalysis Science, Technology* 2018, 8, 6302.
- [52] P. Gartaganis, C. Winkler, Can. J. Chem. 1956, 34, 1457.
- [53] H. Blades, C. Winkler, Can. J. Chem. 1951, 29, 1010.
- [54] T. J. Koller, S. Jin, V. Krol, S. J. Ambach, U. Ranieri, S. Khandarkhaeva, J. Spender, S. McWilliams, F. Trybel, N. Giordano, et al., *Angew. Chem. Int. Ed.* 2024, 63, e202318214.
- [55] E. Horvath-Bordon, R. Riedel, P. McMillan, P. Kroll, G. Miehe, P. van Aken, A. Zerr, P. Hoppe, O. Shebanova, I. McLaren, et al., *Angew. Chem. Int. Ed.* 2007, 46, 1476.
- [56] A. Salamat, K. Woodhead, P. F. McMillan, R. Q. Cabrera, A. Rahman, D. Adriaens, F. Corà, J.-P. Perrillat, *Phys. Rev. B* 2009, 80, 104106.
- [57] M. Sougawa, T. Sumiya, K. Takarabe, Y. Mori, T. Okada, H. Gotou, T. Yagi, D. Yamazaki, N. Tomioka, T. Katsura, et al., *Jpn. J. Appl. Phys.* **2011**, *50*, 095503.
- [58] C. Aldous, Novel van der Waals Compounds in the Nitrogen-Methane Binary System at Room Temperature and High Pressure, Ph.D. thesis 2010.

- [59] I. Nakahata, N. Matsui, Y. Akahama, H. Kawamura, *Chem. Phys. Lett.* **1999**, *302*, 359.
- [60] R. Bini, G. Pratesi, Phys. Rev. B 1997, 55, 14800.
- [61] R. Bini, L. Ulivi, H. J. Jodl, P. R. Salvi, J. Chem. Phys. 1995, 103, 1353.
- [62] M. Hanfland, M. Lorenzen, C. Wassilew-Reul, F. Zontone, *Rev. High Pressure Sci. Techn.* 1998, 7, 787.
- [63] M. Frost, R. T. Howie, P. Dalladay-Simpson, A. F. Goncharov, E. Gregoryanz, *Physical Review B* 2016, 93, 024113.
- [64] Deposition number 2401472 contains the supplementary crystallographic data for this structure. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and FachinformationszentrumKarlsruhe Access Structures service.
- [65] H. Olijnyk, J. Chem. Phys. 1990, 93, 8968.
- [66] M. Bykov, E. Bykova, C. J. Pickard, M. Martinez-Canales, K. Glazyrin, J. S. Smith, A. F. Goncharov, *Phys. Rev. B* 2021, 104, 184105.
- [67] M. I. Eremets, R. J. Hemley, H.-K. Mao, E. Gregoryanz, *Nature* 2001, 411, 170.
- [68] Deposition number 2417277 contains the supplementary crystallographic data for this structure. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and FachinformationszentrumKarlsruhe Access Structures service.
- [69] L. Sun, W. Yi, L. Wang, J. Shu, S. Sinogeikin, Y. Meng, G. Shen, L. Bai, Y. Li, J. Liu, et al., *Chem. Phys. Lett.* **2009**, 473, 72.
- [70] P.M. Bell, H.K. Mao, R.J. Hemley, *Physica B+C* 1986, 139, 16.
- [71] D. Tomasino, M. Kim, J. Smith, C.-S. Yoo, Phys. Rev. Lett. 2014, 113, 205502.
- [72] F. Zahariev, S. V. Dudiy, J. Hooper, F. Zhang, T. K. Woo, *Phys. Rev. Lett.* **2006**, *97*, 155503.
- [73] M. Kirsz, C.G. Pruteanu, P.I.C. Cooke, G.J. Ackland, Understanding solid nitrogen through machine learning simulation 2024.
- [74] S. S. Lobanov, P.-N. Chen, X.-J. Chen, C.-S. Zha, K. D. Litasov, H.-K. Mao, A. F. Goncharov, *Nat. Commun.* 2013, 4, 2446.
- [75] E. Stavrou, S. Lobanov, H. Dong, A. R. Oganov, V. B. Prakapenka, Z. Konôpková, A. F. Goncharov, *Chem. Mater.* 2016, 28, 6925.
- [76] A. A. da Costa, C. Geraldes, J. Teixeira-Dias, J. Raman Spectrosc. 1982, 13, 56.
- [77] J. Durig, S. Bush, F. Baglin, J. Chem. Phys. 1968, 49, 2106.
- [78] E. Sato, H. Martinho, *Biomed. Optics Express* 2018, 9, 1728.
 [79] M. Cable, S. Hörst, C. He, A. Stockton, M. Mora, M. Tolbert,
- [79] M. Cable, S. Holst, C. He, A. Stockton, M. Mola, M. Holbert, M. Smith, P. Willis, *Earth Planet. Sci. Lett.* 2014, 403, 99.
- [80] F. Raulin, C. Brassé, O. Poch, P. Coll, Chem. Soc. Rev. 2012, 41, 5380.
- [81] F. Datchi, S. Ninet, M. Gauthier, A.M. Saitta, B. Canny, F. Decremps, *Phys. Rev. B* 2006, *73*, 174111.
- [82] A. Serovaiskii, V. Kutcherov, Sci. Rep. 2020, 10, 4559.
- [83] M. Wang, M. Pena-Alvarez, R. Howie, E. Gregoryanz, Revisiting the Phase Diagram of Methane 2024, submitted.

Manuscript received: November 21, 2024

Accepted manuscript online: February 19, 2025 Version of record online: **D**.

Angew. Chem. Int. Ed. 2025, e202422710 (7 of 7)



Research Article

High-Pressure Chemistry

H. A. Shuttleworth, M. A. Kuzovnikov,
L. J. Conway, H. Hu, J. Yan, S. Gallego-Parra, I. Osmond, T. Marqueño,
M. Hanfland, D. Laniel, E. Gregoryanz,
A. Hermann,* M. Peña-Alvarez,*
R. T. Howie* _________ e202422710

Pressure-Driven Reactivity in Dense Methane-Nitrogen Mixtures



 CH_4 and N_2 are abundant molecules in our solar system and are the primary constituents of Titan's atmosphere. Under extreme pressures and temperatures within a diamond anvil cell, CH_4 and N_2 are demonstrated to react to form a range of compounds, dependent on the conditions and initial concentrations. This provides valuable insight into the complexity of the C–N–H ternary system under planetary conditions.